

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

010180.00012

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

10/049274

INTERNATIONAL APPLICATION NO.
PCT/GB99/02629INTERNATIONAL FILING DATE
August 10, 1999

PRIORITY DATE CLAIMED

TITLE OF INVENTION

ANTIBACTERIAL AGENTS

APPLICANT(S) FOR DO/EO/US

Michael George HUNTER, Paul Raymond BECKETT, John Martin CLEMENTS, Mark WHITTAKER, Stephen John DAVIES, Lisa Marie PRATT, Zoe Marie SPAVOLD, Steven LAUNCHBURY


Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☒ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☒ A copy of the International Search Report (PCT/ISA/210).

Items 13 to 20 below concern document(s) or information included:

13. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☐ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☐ Certificate of Mailing by Express Mail
23. ☒ Other items or information:


PCT REQUEST (5pp.); PCT/IPEA/416 (1 p.); Copy of WO 01/10835 published Feb. 15, 2001 w/PCT/ISA/210: Specification (40 pp.), Claims 1-5 (4 pp.), Abstract (1 p.) PCT/IPEA/409 (5pp.) w/Claims 1-4 (Amended Sheets: 7); PCT/IB/308; PCT/IB/332

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 10/049274)		INTERNATIONAL APPLICATION NO. PCT/GB99/02629		ATTORNEY'S DOCKET NUMBER 010180.00012	
24. The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (I) - (5)) : <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).				\$890.00	
				\$0.00	
CLAIMS		NUMBER FILED		NUMBER EXTRA	
RATE					
Total claims	4 - 20 =	0	x \$18.00	\$0.00	
Independent claims	2 - 3 =	0	x \$84.00	\$0.00	
Multiple Dependent Claims (check if applicable).				<input type="checkbox"/>	\$0.00
TOTAL OF ABOVE CALCULATIONS =				\$890.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27). The fees indicated above are reduced by 1/2.				\$0.00	
SUBTOTAL =				\$890.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).				\$0.00	
TOTAL NATIONAL FEE =				\$890.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable).				<input type="checkbox"/>	\$0.00
TOTAL FEES ENCLOSED =				\$890.00	
				Amount to be:	\$
				refunded	
				charged	\$
a. <input type="checkbox"/> A check in the amount of _____ to cover the above fees is enclosed.					
b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. 19-0733 in the amount of \$890.00 to cover the above fees. A duplicate copy of this sheet is enclosed.					
c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 19-0733 . A duplicate copy of this sheet is enclosed.					
d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO:					
BANNER & WITCOFF, LTD. 1001 G Street, N.W., 11th Floor Washington, D.C 20001			 SIGNATURE Susan A. WOLFFE NAME 33,568 REGISTRATION NUMBER February 11, 2002 DATE		

REMARKS

The amendment to the specification is made in accordance with 35 U.S.C. 119, 37
C.F.R. 1.78 and 37 C.F.R. 1.55. Entry is requested.

Respectfully submitted,


Susan A. Wolfe
Reg. No. 33,568

February 11, 2002
BANNER & WITCOFF, LTD.
Eleventh Floor
1001 G Street, N.W.
Washington, D.C. 20001-4597
(202) 508-9100

10/04 9274

Rec'd PCT/PTO 11 FEB 2002

WO 01/10835

PCT/GB99/02629

1

Antibacterial Agents

This invention relates to the use of N-formyl hydroxylamine derivatives as antibacterial agents, to a novel class of such compounds, and to pharmaceutical and veterinary compositions comprising such compounds.

Background to the Invention

In general, bacterial pathogens are classified as either Gram-positive or Gram-negative. Many antibacterial agents (including antibiotics) are specific against one or other Gram-class of pathogens. Antibacterial agents effective against both Gram-positive and Gram-negative pathogens are therefore generally regarded as having broad spectrum activity.

Many classes of antibacterial agents are known, including the penicillins and cephalosporins, tetracyclines, sulfonamides, monobactams, fluoroquinolones and quinolones, aminoglycosides, glycopeptides, macrolides, polymyxins, lincosamides, trimethoprim and chloramphenicol. The fundamental mechanisms of action of these antibacterial classes vary.

Bacterial resistance to many known antibacterials is a growing problem. Accordingly there is a continuing need in the art for alternative antibacterial agents, especially those which have mechanisms of action fundamentally different from the known classes.

Amongst the Gram-positive pathogens, such as Staphylococci, Streptococci, Mycobacteria and Enterococci, resistant strains have evolved/arisen which makes them particularly difficult to eradicate. Examples of such strains are methicillin resistant *Staphylococcus aureus* (MRSA), methicillin resistant coagulase negative Staphylococci (MRCNS), penicillin resistant *Streptococcus pneumoniae* and multiply resistant *Enterococcus faecium*.

Pathogenic bacteria are often resistant to the aminoglycoside, β -lactam (penicillins and cephalosporins), and chloramphenicol types of antibiotic. This resistance involves the enzymatic inactivation of the antibiotic by hydrolysis or by formation of inactive derivatives. The β -lactam (penicillin and cephalosporin) family of antibiotics are characterised by the presence of a β -lactam ring structure. Resistance to this family of antibiotics in clinical isolates is most commonly due to the production of a "penicillinase" (β -lactamase) enzyme by the resistant bacterium which hydrolyses the β -lactam ring thus eliminating its antibacterial activity.

Recently there has been an emergence of vancomycin-resistant strains of enterococci (Woodford N. 1998 Glycopeptide-resistant enterococci: a decade of experience. *Journal of Medical Microbiology*. 47(10):849-62). Vancomycin-resistant enterococci are particularly hazardous in that they are frequent causes of hospital based infections and are inherently resistant to most antibiotics. Vancomycin works by binding to the terminal D-Ala-D-Ala residues of the cell wall peptidoglycan precursor. The high-level resistance to vancomycin is known as VanA and is conferred by a genes located on a transposable element which alter the terminal residues to D-Ala-D-lac thus reducing the affinity for vancomycin.

In view of the rapid emergence of multidrug-resistant bacteria, the development of antibacterial agents with novel modes of action that are effective against the growing number of resistant bacteria, particularly the vancomycin resistant enterococci and β -lactam antibiotic-resistant bacteria, such as methicillin-resistant *Staphylococcus aureus*, is of utmost importance.

Brief Description of the Invention

This invention is based on the finding that certain N-formyl hydroxylamine derivatives have antibacterial activity, and makes available a new class of antibacterial agents. The inventors have found that the compounds with which this invention is concerned are antibacterial with respect to a range of Gram-positive and Gram-negative organisms.

Although it may be of interest to establish the mechanism of action of the compounds with which the invention is concerned, it is their ability to inhibit bacterial growth that makes them useful. However, it is presently believed that their antibacterial activity is due, at least in part, to intracellular inhibition of bacterial polypeptide deformylase (PDF; EC 3.5.1.31).

All ribosome-mediated synthesis of proteins starts with a methionine residue. In prokaryotes the methionyl moiety carried by the initiator tRNA is N-formylated prior to its incorporation into a polypeptide. Consequently, N-formylmethionine is always present at the N-terminus of a nascent bacterial polypeptide. However, most mature proteins do not retain the N-formyl group or the terminal methionine residue. Deformylation is required prior to methionine removal, since methionine aminopeptidase does not recognise peptides with an N-terminal formylmethionine residue (Solbiati et al., J. Mol. Biol. 290:607-614, 1999). Deformylation is, therefore, a crucial step in bacterial protein biosynthesis and the enzyme responsible, PDF, is essential for normal bacterial growth. Although the gene encoding PDF (*def*) is present in all pathogenic bacteria for which sequences are known (Meinzel et al., J. Mol. Biol. 266:939-49, 1997), it has no eukaryotic counterpart, making it an attractive target for antibacterial chemotherapy.

The isolation and characterisation of PDF has been facilitated by an understanding of the importance of the metal ion in the active site (Groche et al., Biophys. Biochem. Res. Commun., 246:324-6, 1998). The Fe^{2+} form is highly active *in vivo* but is unstable when isolated due to oxidative degradation (Rajagopalan et al., J. Biol. Chem. 273:22305-10, 1998). The Ni^{2+} form of the enzyme has specific activity comparable with the ferrous enzyme but is oxygen-insensitive (Ragusa et al., J. Mol. Biol. 1998, 280:515-23, 1998). The Zn^{2+} enzyme is also stable but is almost devoid of catalytic activity (Rajagopalan et al., J. Am. Chem. Soc. 119:12418-12419, 1997).

Several X-ray crystal structures and NMR structures of *E. coli* PDF, with or without bound inhibitors, have been published (Chan et al., Biochemistry 36:13904-9, 1997; Becker et al., Nature Struct. Biol. 5:1053-8, 1998; Becker et al., J. Biol. Chem. 273:11413-6, 1998; Hao et al., Biochemistry, 38:4712-9, 1999; Dardel et al., J. Mol. Biol. 280:501-13, 1998; O'Connell et al., J. Biomol. NMR, 13:311-24, 1999), indicating similarities in active site geometry to metalloproteinases such as thermolysin and the metzincins.

Recently the substrate specificity of PDF has been extensively studied (Ragusa et al., J. Mol. Biol. 289:1445-57, 1999; Hu et al., Biochemistry 38:643-50, 1999; Meinnel et al., Biochemistry, 38:4287-95, 1999). These authors conclude that an unbranched hydrophobic chain is preferred at P1', while a wide variety of P2' substituents are acceptable and an aromatic substituent may be advantageous at the P3' position. There have also been reports that small peptidic compounds containing an H-phosphonate (Hu et al., Bioorg. Med. Chem. Lett., 8:2479-82, 1998) or thiol (Meinnel et al., Biochemistry, 38:4287-95, 1999) metal binding group are micromolar inhibitors of PDF. Peptide aldehydes such as calpeptin (N-Cbz-Leu-norleucinal) have also been shown to inhibit PDF (Durand et al., Arch. Biochem. Biophys., 367:297-302, 1999). However, the identity of the metal binding group and its spacing from the rest of the molecule ("recognition fragment") has not been studied extensively. Furthermore, non-peptidic PDF inhibitors, which may be desirable from the point of view of bacterial cell wall permeability or oral bioavailability in the host species, have not been identified.

Related Prior Art

Certain N-formyl hydroxylamine derivatives have previously been claimed in the patent publications listed below, although very few examples of such compounds have been specifically made and described:


EP-B-0236872	(Roche)
WO 92/09563	(Glycomed)

WO 92/04735	(Syntex)
WO 95/19965	(Glycomed)
WO 95/22966	(Sanofi Winthrop)
WO 95/33709	(Roche)
WO 96/23791	(Syntex)
WO 96/16027	(Syntex/Agouron)
WO 97/03783	(British Biotech)
WO 97/18207	(DuPont Merck)
WO 98/38179	(GlaxoWellcome)
WO 98/47863	(Labs Jaques Logeais)



The pharmaceutical utility ascribed to the N-formyl hydroxylamine derivatives in those publications is the ability to inhibit matrix metalloproteinases (MMPs) and in some cases release of tumour necrosis factor (TNF), and hence the treatment of diseases or conditions mediated by those enzymes, such as cancer and rheumatoid arthritis. That prior art does not disclose or imply that N-formyl hydroxylamine derivatives have antibacterial activity.

In addition to these, US-A-4,738,803 (Roques et al.) also discloses N-formyl hydroxylamine derivatives, however, these compounds are disclosed as enkephalinase inhibitors and are proposed for use as antidepressants and hypotensive agents. Also, WO 97/38705 (Bristol-Myers Squibb) discloses certain N-formyl hydroxylamine derivatives as enkephalinase and angiotensin converting enzyme inhibitors. This prior art does not disclose or imply that N-formyl hydroxylamine derivatives have antibacterial activity either.

Our copending International Patent Application No. PCT/GB99/0386 describes and claims, *inter alia*, the use of a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof in the preparation of an antibacterial composition:



wherein R_2 represents a substituted or unsubstituted C_1 - C_6 alkyl, cycloalkyl(C_1 - C_6 alkyl)-, or aryl(C_1 - C_6 alkyl)- group, and A represents a group of formula (IA), or (IB):



wherein R_4 represents the side chain of a natural or non-natural alpha amino acid, and R_5 and R_6 are each independently hydrogen or C_1 - C_6 alkyl, heterocyclic or aryl(C_1 - C_6 alkyl)-, or R_5 and R_6 when taken together with the nitrogen atom to which they are attached form an optionally substituted saturated heterocyclic ring of 3 to 8 atoms which ring is optionally fused to a carbocyclic or second heterocyclic ring.

Detailed description of the invention

The present invention provides additional members of the class of compounds disclosed in PCT/GB99/00386, but which were not specifically identified or exemplified therein. As members of the class disclosed in PCT/GB99/00386, the present compounds are antibacterially active, and their mechanism of action is presently believed to be due at least in part to their ability to inhibit bacterial peptide deformylases.

Accordingly, the present invention provides a compound of formula (I) as defined above, selected from the group consisting of:

N-[3S-(4-benzylpiperidine-1-carbonyl)-2,2-dimethyl-propyl]-3-cyclopentyl-2R-
[(formyl-hydroxy-amino)-methyl]-propionamide,

N-[2R-(4-benzyl-piperidine-1-carbonyl)-hexyl]-N-hydroxy-formamide,

N-hydroxy-N-[2R-(2-methyl-piperidine-1-carbonyl)-hexyl]-formamide,

N-hydroxy-N-[2R-(piperidine-1-carbonyl)-hexyl]-formamide,

N-hydroxy-N-[2R-(piperazine-1-carbonyl)-hexyl]-formamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid pyrrolidin-1-ylamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid methyl-(1-methyl-
piperidin-4-yl)-amide,

N-[2R-(azepane-1-carbonyl)-hexyl]-N-hydroxy-formamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (4-methyl-piperazin-1-yl)-
amide,

2R-[(formyl-hydroxy-amino)-methy]-hexanoic acid diisopropylamide,

1-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl}-piperidine-3-carboxylic acid
ethyl ester,

4-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl}-piperazine-1-carboxylic acid
ethyl ester,

4-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl}-1,1-dimethyl-piperazinium

iodide,

2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid [2,2-dimethyl-1S-(piperidine-1-carbonyl)-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid
[1S-(3,4-dihydro-1*H*-isoquinoline-2-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-4-hydroxy-piperidine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-piperazine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (3-benzylsulfanyl-1S-dimethylcarbamoyl-propyl)-amide,

3S-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoylamino}-*N,N*-dimethyl-succinamic acid benzyl ester,

4S-dimethylcarbamoyl-4-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl-amino}-butyric acid benzyl ester,

(5S-dimethylcarbamoyl-5-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl-amino}-pentyl)-dimethyl-ammonium chloride,

2R-[(formyl-hydroxy-amino)-methyl]-butyric acid (1-carbamoyl-2,2-dimethyl-propyl) amide,

2-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (1-carbamoyl-2,2-dimethyl-propyl) amide,

2R-[formyl-hydroxy-amino)-methyl]-hexanoic acid (1-dimethyl-carbamoyl-4-guanidinobutyl)-amide,

2R-[2-(4-chlorophenyl)-3-(formyl-hydroxy-amino)-propionylamino]-2S-3,3,N,N-tetramethyl-butylamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [2(3,4-dihydroxy-phenyl)-ethyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [2(4-hydroxyphenyl)-ethyl]-amide,

and pharmaceutically and veterinarily acceptable salts, hydrates and solvates thereof.

According to other aspects of the invention, there is provided (a) the use of a compound specifically named immediately above, or a pharmaceutically or veterinarily acceptable salt solvate or hydrate thereof, in the preparation of an antibacterial composition; (b) a method for the treatment of bacterial infections in humans and non-human mammals, which comprises administering to a subject suffering such infection an antibacterially effective dose of a compound specifically named immediately above, or a pharmaceutically or veterinarily acceptable salt solvate or hydrate thereof; (c) a method for the treatment of bacterial contamination by applying an antibacterially effective amount of a compound specifically named immediately above, or a pharmaceutically or veterinarily acceptable salt solvate or hydrate thereof, to the site of contamination; and (d) a pharmaceutical or veterinary composition comprising a compound specifically named immediately above, or a pharmaceutically or veterinarily acceptable salt solvate or hydrate thereof, together with a pharmaceutically or veterinarily acceptable carrier.

Of the compounds of the invention, the following are presently especially preferred

for their potency as antibacterial agents:

N-[3S-(4-benzylpiperidine-1-carbonyl)-2,2-dimethyl-propyl]-3-cyclopentyl-2R-[(formyl-hydroxy-amino)-methyl]-propionamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid methyl-(1-methyl-piperidin-4-yl)-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-4-hydroxy-piperidine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-piperazine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (3-benzylsulfanyl-1S-dimethylcarbamoyl-propyl)-amide, and

2-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (1-carbamoyl-2,2-dimethyl-propyl) amide.

On the hypothesis that the compounds (I) act by inhibition of intracellular PDF, the most potent antibacterial effect may be achieved by using compounds which efficiently pass through the bacterial cell wall. Thus, compounds which are highly active as inhibitors of PDF in vitro and which penetrate bacterial cells are preferred for use in accordance with the invention. It is to be expected that the antibacterial potency of compounds which are potent inhibitors of the PDF enzyme in vitro, but are poorly cell penetrant, may be improved by their use in the form of a prodrug, ie a structurally modified analogue which is converted to the parent molecule of formula (I), for example by enzymic action, after it has passed through the bacterial cell wall.

Salts of the compounds of the invention include physiologically acceptable acid addition salts for example hydrochlorides, hydrobromides, sulphates, methane sulphonates, p-toluenesulphonates, phosphates, acetates, citrates, succinates, lactates, tartrates, fumarates and maleates. Salts may also be formed with bases, for example sodium, potassium, magnesium, and calcium salts.

Compositions with which the invention is concerned may be prepared for administration by any route consistent with the pharmacokinetic properties of the active ingredient(s).

Orally administrable compositions may be in the form of tablets, capsules, powders, granules, lozenges, liquid or gel preparations, such as oral, topical, or sterile parenteral solutions or suspensions. Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinyl-pyrrolidone; fillers for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricant, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants for example potato starch, or acceptable wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, glucose syrup, gelatin hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

For topical application to the skin, the active ingredient(s) may be made up into a cream, lotion or ointment. Cream or ointment formulations which may be used for the drug are conventional formulations well known in the art, for example as described in standard textbooks of pharmaceuticals such as the British Pharmacopoeia.

The active ingredient(s) may also be administered parenterally in a sterile medium. Depending on the vehicle and concentration used, the drug can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle. Intra-venous infusion is another route of administration for the compounds used in accordance with the invention.

Safe and effective dosages for different classes of patient and for different disease states will be determined by clinical trial as is required in the art. It will be understood that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The following Examples describe the preparation of the compounds of the invention. In the Examples, ^1H and ^{13}C NMR spectra were recorded using a Bruker DPX 250 spectrometer at 250.1 and 62.9MHz, respectively. Mass spectra were obtained using a Perkin Elmer Sciex API 165 spectrometer using both positive and negative ionisation modes. Infra-red spectra were recorded on a Perkin Elmer PE 1600 FTIR spectrometer. The following abbreviations have been used throughout:

DIAD	Diisopropylazodicarboxylate
DIPEA	Diisopropylethylamine
DMF	N,N-Dimethylformamide

WO 01/10835


PCT/GB99/02629

13

EDC	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
HOAt	1-Hydroxy-7-aza-benzotriazole
HOBt	1-Hydroxybenzotriazole
LRMS	Low resolution mass spectrometry
THF	Tetrahydrofuran


Example 1

N-[3S-(4-Benzylpiperidine-1-carbonyl)-2,2-dimethyl-propyl]-3-cyclopentyl-2R-[(formyl-hydroxy-amino)-methyl]-propionamide



The title compound was prepared as detailed below (see also Scheme 1)

Scheme 1



Reagents and Conditions: A: TiCl_4 , trioxane, CH_2Cl_2 ; B: H_2O_2 , LiOH ; C: H_2NOBn , WSC, $\text{THF}/\text{H}_2\text{O}$; D: Ph_3P , DIAD, THF ; E: LiOH , $\text{THF}/\text{MeOH}/\text{H}_2\text{O}$; F: formic acetic anhydride, NEt_3 , THF ; G: H-Tle-amide, EDCI, HOAt, DMF; H: Pd/C , H_2 , MeOH .

Step A: 4S-Benzyl-3-[3-cyclopentyl-2R-hydroxymethyl-propionyl]-oxazolidin-2-one

To a stirred, cooled (0°C) solution of 4S-benzyl-(3-cyclopentyl-propionyl)-oxazolidin-2-one (21 g, 69.8 mmol) in dichloromethane (350 ml) was added a solution of titanium tetrachloride (1M in dichloromethane, 73.25 ml, 73.2 mmol), dropwise. The resulting yellowish slurry was stirred for 10 minutes at 0°C , and then DIPEA (13.37 ml, 76.7 ml) was added dropwise to furnish a dark-red solution. The stirring was maintained for 1 h at 0°C , and then a solution of s-trioxane (7.53 g, 83.7 mmol), in dichloromethane (70 ml) was added dropwise followed by the addition of a solution of titanium tetrachloride (1M in dichloromethane, 73.25 ml, 73.2 mmol). The reaction mixture was then stirred

for 4 h at 0 °C. Saturated aqueous ammonium chloride (250 ml) was added to the reaction mixture and the aqueous layer was extracted with additional dichloromethane (2x300 ml). The combined organic layers were washed with water (150 ml) and with brine (80 ml), dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to yield a yellow solid which on trituration with diethyl ether furnished a white solid (16.57 g, 72%). ¹H-NMR; δ(CDCl₃), 7.38-7.22 (5H, m), 4.70 (1H, m), 4.22-4.18 (2H, m), 3.99 (1H, m), 3.96-3.75 (2H, m), 3.31 (1H, dd, J = 13.4 & 3.3 Hz), 2.82 (1H, dd, J = 13.4 & 9.4 Hz), 2.24 (1H, dd, J = 8.3 & 4.5 Hz), 2.81-1.30 (4H, m) and 1.13 (1H, m); ¹³C-NMR; δ(CDCl₃), 176.3, 153.6, 135.2, 129.5, 129.0, 127.4, 66.2, 64.2, 55.7, 44.8, 37.9, 37.8, 34.6, 33.0, 32.4 and 25.1.

Step B: 3-Cyclopentyl-2R-hydroxymethyl-propionic acid

To a stirred, cooled (0 °C) solution of 4S-Benzyl-3-[3-cyclopentyl-2R-hydroxymethyl-propionyl]-oxazolidin-2-one (16.05 g, 48.5 mmol) in THF-water (4:1, 250 ml) was added 27.5% aqueous hydrogen peroxide (24 ml, 194 mmol), followed by lithium hydroxide monohydrate (4.07 g, 97 mmol) in water (50 ml). After the reaction was complete (30 min), THF was removed *in vacuo*. The aqueous layer was extracted with dichloromethane (3x100 ml) and acidified to pH 2 with 4M hydrochloric acid. The aqueous layer was extracted with diethyl ether (2x150 ml). The combined organic layers were washed with brine (60 ml), dried over anhydrous magnesium sulfate and filtered. The solvent was removed *in vacuo* to afford a yellow oil which was further purified by column chromatography (25% ethyl acetate in hexanes to 100% ethyl acetate) to furnish the title compound as an oil (8.3 g, quant.). ¹H-NMR; δ (CDCl₃), 6.60-5.90 (1H, br s), 3.80-3.78 (2H, m), 2.67 (1H, m), 1.98-1.40 (9H, m) and 1.20-0.98 (2H, m). ¹³C-NMR; δ(CDCl₃), 181.0, 63.2, 46.9, 37.8, 34.5, 32.7, 32.6, 25.1 and 25.1.

Step C: N-Benzoyloxy-3-cyclopentyl-2R-hydroxymethyl-propionamide

To a stirred, cooled (0 °C) mixture of 3-cyclopentyl-2R-hydroxymethyl-propionic acid (1.1 g, 6.4 mmol) in THF-water (4:1, 30 ml), was added O-benzylhydroxylamine. The

WO 01/10835

PCT/GB99/02629

16

pH of the resulting solution was adjusted to 4.5 by addition of 1M hydrochloric acid, and then EDC (1.84 g, 9.6 mmol) was added in one portion. The resulting solution was stirred for 2.5 h at room temperature while controlling pH at 4.5 by addition of 1M hydrochloric acid. After removal of the THF, the aqueous layer was extracted with ethyl acetate (3x40 ml) and the combined organic layers were washed with 10% citric acid (3x15 ml), 5% sodium hydrogen carbonate and dried over anhydrous magnesium sulfate. The solvent was removed *in vacuo* to afford the title compound as a colourless crystalline solid (1.18 g, 67%). This compound was then used without any further purification. ¹H-NMR; δ (CDCl₃), 8.14 (1H, br s), 7.40-7.34 (5H, m), 4.94 (2H, br s), 3.76-3.66 (2H, m), 1.79-1.47 (11H, m) and 1.17-0.97 (2H, m). LRMS: +ve ion 278 [M+H], 555 [2M+H].

Step D: N-Benzyloxy-3R-cyclopentylmethyl-azetidin-2-one

To a stirred, cooled (0 °C) solution of *N*-Benzyloxy-3-cyclopentyl-2R-hydroxymethyl-propionamide (8.63 g, 31.1 mmol) and triphenylphosphine (9 g, 34.2 mmol) in dry THF (320 ml) was added DIAD (6.12 ml, 31.1 mmol), dropwise. The resulting solution was stirred at room temperature overnight. After removal of THF *in vacuo*, the residue was purified by column chromatography (hexanes:ethyl acetate, 5:1 to 3:1) to give the desired product as a white solid (6.7 g, 83%). ¹H-NMR; δ (CDCl₃), 7.76-7.39 (5H, m), 4.94 (2H, br s), 3.36 (1H, m), 2.96-2.80 (2H, m), 1.89-1.38 (9H, m) and 1.18-0.98 (2H, m). ¹³C-NMR; δ (CDCl₃), 167.7, 129.6, 129.3, 129.0, 78.1, 52.5, 45.1, 39.1, 35.2, 33.1, 32.9, 25.5 and 25.3. LRMS: +ve ion 260 [M+H], 519 [2M+H].

Step E: 2R-(Benzyloxyamino-methyl)-3-cyclopentyl-propionic acid

To a stirred, cooled (0 °C) solution of *N*-Benzyloxy-3R-cyclopentylmethyl-azetidin-2-one (6.7 g, 25.8 mmol) in THF-methanol (3:1, 100 ml) was added lithium hydroxide monohydrate (1.3 g, 31.0 mmol) in water (25 ml). The reaction mixture was stirred and allowed to warm to room temperature overnight. The solvent was removed *in vacuo* and the aqueous layer was extracted with diethyl ether, then acidified to pH 2 by

addition of 4M hydrochloric acid. The aqueous layer was extracted with diethyl ether (3x40 ml), and the combined organic layers were washed with brine, dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to give the title compound as white crystals (6.02 g, 84%). ¹H-NMR; δ (CDCl₃), 7.68-7.30 (5H, m), 4.78-4.68 (2H, m), 3.12-3.10 (2H, d, J = 6.9 Hz), 2.76 (1H, m), 1.91-1.39 (11H, m), 1.20-1.00 (2H, m). ¹³C-NMR; δ (CDCl₃), 180.1, 137.7, 129.0, 128.9, 128.5, 78.0, 53.9, 42.9, 38.3, 36.6, 33.1, 33.0, 25.5. LRMS: -ve ion 276 [M-H], 553 [2M-H].

Step F: 2R-[(Benzyloxy-formyl-amino)-methyl]-3-cyclopentyl-propionic acid

To a stirred, cooled (0 °C) solution of 2R-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid (3.79 g, 13.7 mmol) in THF (20 ml) was added formic acetic anhydride (3.01 g, 34.2 mmol) and triethylamine (5.72 ml, 41.0 mmol). The reaction mixture was stirred for 1 h at 0 °C and 45 min at room temperature. The solvent was removed *in vacuo* and the mixture was purified by flash chromatography (hexanes: ethyl acetate, 1:1) to give the title compound as a yellow oil (3.04 g, 73%). LRMS: -ve ion 304 [M-H], -ve ion 609 [2M-H].

Step G: 2R-[(Benzyloxy-formyl-amino)-methyl]-N-[1S-(4-benzyl-piperidine-1-carbonyl)-2,2-dimethyl-propyl]-3-cyclopentyl-propionamide

To a stirred, cooled (0 °C) solution of 2R-[(Benzyloxy-formyl-amino)-methyl] -3-cyclopentyl-propionic acid (396 mg, 1.3 mmol) and 2S-Amino-1-(4-benzyl-piperidin-1-yl)-3,3-dimethyl-butan-1-one (see below) in DMF (5 ml), were added EDC (274 mg, 1.43 mmol) and HOAt (8.8 mg, 0.065 mmol). The reaction mixture was stirred overnight at room temperature. DMF was removed *in vacuo* to furnish a yellow oil, which was dissolved in ethyl acetate. The organic layer was then washed with 1M hydrochloric acid (2x5 ml) and water (5 ml). The aqueous layer was re- extracted with ethyl acetate. The combined organic layers were washed with brine, dried over anhydrous magnesium sulfate, filtered and


the solvent was removed *in vacuo* to furnish a white foam (660 mg, 88%) which was used in the next step without any purification.

Step H: *N*-[1*S*-(4-benzylpiperidine-1-carbonyl)-2,2-dimethyl-propyl]-3-cyclopentyl-2*R*-[(formyl-hydroxy-amino)-methyl]-propionamide

To a stirred solution of the 2*R*-[(benzyloxy-formyl-amino)-methyl]-*N*-[1*S*-(4-benzylpiperidine-1-carbonyl)-2,2-dimethyl-propyl]-3-cyclopentyl-propionamide (655 mg, 1.14 mmol) in Methanol (8 ml) under an argon atmosphere was added Pd/C (70 mg). Hydrogen gas was bubbled through the suspension for 30 min. The reaction mixture was then filtered through celite and the solvent was removed *in vacuo* to afford a pale pink solid (522 mg, 95%). ¹H-NMR; δ (CDCl₃, rotamers), 8.40 (0.4H, m), 7.83 (0.6H, m), 7.34-7.09 (5H, m), 6.55 (1H, m), 4.90 (1H, m), 4.57 (1H, m), 4.11-3.99 (1.5H, m), 3.85-3.77 (0.8H, m), 3.63-3.59 (0.7H, m), 3.51-3.47 (0.6H, m), 3.08-2.95 (1.2H, m), 2.88-2.62 (1.2H, m), 2.57-2.49 (3H, m), 1.89-0.90 (25H, m). LRMS: +ve ion 508 [M+Na], -ve ion 484 [M-H].

Preparation of 2*S*-Amino-1-(4-benzyl-piperidin-1-yl)-3,3-dimethyl-butan- 1-one (see Scheme 2)

Scheme 2



Reagents and conditions: A. NEt₃, N-(benzyloxycarbonyloxy)-succinimide, MeOH; B. EDCI, HOAt, DMF; C. cyclohexene, Pd/C, EtOH

Step A: 2S-Benzyloxycarbonylamino-3,3-dimethyl-butyric acid

To a suspension of L-*tert*-leucine (11.88 g, 90.7 mmol) in methanol (200 ml) were added triethylamine (26.56 ml, 190 mmol) and *N*-(benzyloxycarbonyl- oxy)-succinimide (24.88 g, 99.8 mmol). The reaction mixture was stirred at room temperature for 14 h. Methanol was removed *in vacuo* to afford a viscous pale yellow oil, which was dissolved in ethyl acetate (100 ml). The organic layer was washed with 1M hydrochloric acid (15 ml) and brine, dried over anhydrous magnesium sulfate and filtered. The solvent was removed *in vacuo* to furnish the title compound as an oil (24 g, quant.). ¹H-NMR; δ(CDCl₃), 7.43-7.36 (5H, m), 5.36 (1H, d, J = 9.4 Hz), 5.12 (2H, br s), 4.20 (1H, d, J = 9.6 Hz) and 1.02 (9H, s). LRMS: +ve ion 266 [M+H], -ve ion 264 [M-H], 529 [2M-H].

Step B: 2S-[1-(4-Benzyl-piperidine-1-carbonyl)-2,2-dimethyl-propyl]-carbamic acid benzyl ester


To a solution of 2S-Benzyloxycarbonylamino-3,3-dimethyl-butyric acid (923 mg, 3.48 mmol) and 4-benzyl piperidine (735 μl, 4.18 mmol) in DMF (16 ml) were added EDC (734 mg, 3.83 mmol) and HOAt (10 mg, 0.07 mmol). The reaction mixture was stirred for 14 h at room temperature. DMF was removed *in vacuo* and the crude residue was dissolved in ethyl acetate. The organic layer was washed with 1M hydrochloric acid (2x10 ml), water (10 ml), brine (10 ml), dried over anhydrous magnesium sulfate and filtered. Removal of the solvent *in vacuo* and purification by column chromatography (hexanes:ethyl acetate, 5:1) provided the desired amide (784 mg, 54%). ¹H-NMR; δ(CDCl₃), 7.36-7.14 (10H, m), 5.65 (1H, m), 5.17-5.05 (2H, m), 4.70-4.49 (2H, m), 2.96 (1H, m), 2.57-2.47 (2H, m), 1.90-1.59 (2H, m) and 1.38-0.87 (14H, m). LRMS: +ve ion 423 [M+H].

Step C: 2S-Amino-1-(4-benzyl-piperidin-1-yl)-3,3-dimethyl-butan-1-one

To a stirred solution of 2S-[1-(4-Benzyl-piperidine-1-carbonyl)-2,2-dimethyl-propyl-carbamic acid benzyl ester (784 mg, 1.86 mmol) in ethanol (4 ml) was added 10% palladium on charcoal (70 mg) and cyclohexene (380 μ l, 3.71 mmol). The suspension was warmed to 75 °C for 1.5 h. The reaction mixture was filtered through celite and the solvent was removed *in vacuo* to afford quantitatively the title compound as a viscous oil. $^1\text{H-NMR}$; $\delta(\text{CDCl}_3)$, 7.32-7.12 (5H, m), 4.69 (1H, m), 4.01 (1H, m), 3.53 (1H, m), 2.86 (1H, m), 2.63-2.45 (3H, m), 1.80-1.63 (3H, m), 1.30-1.08 (3H, m), 0.99 (4.5H, m) and 0.94 (4.5H, m). LRMS: +ve ion 289 [M+H].

Examples 2-12

The compounds of Examples 2-12 (Table 1) were prepared in array format using the generic procedure outlined below (see also Scheme 3).

Scheme 3

Reagents and conditions: A. piperidine, HCHO, EtOH, 80°C, o/n; B. $^t\text{BuCOCl}$, Et_3N then 3-lithio-4-benzyl-5,5-dimethyl-oxazolidin-2-one; C. H_2NOBzl , room temp., o/n then pTsOH , EtOAc ; D. LiOH , aq THF, 0°C; E. formic acetic anhydride, Et_3N , THF; F. PfpOH , EDC , HOBT , THF; G. Amine; H. cyclohexene, Pd/C , EtOH.

Analytical HPLC was performed on a Beckman System Gold, using Waters Nova Pak C18 column (150 mm, 3.9 mm) with 20 to 90 % solvent B gradient (1 ml/min) as the mobile phase. [Solvent A: 0.05% TFA in 10% water 90% methanol; Solvent B: 0.05% TFA in 10% methanol 90%], detection wavelength at 230 nm. Preparative HPLC was performed on a Gilson autoprep instrument using a C18 Waters delta prep-pak cartridge (15 μ m, 300 A, 25 mm, 10 mm) with 20 to 90 % solvent B gradient (6 ml/min) as the mobile phase. [Solvent A water; Solvent B: methanol], UV detection was at 230 nm.

Step A: 2-Butyl acrylic acid

To a solution of n-butylnalonic acid (17.2 g, 107 mmol) in ethanol (200 ml) was added piperidine (12.76 ml, 129 mmol) and 37% aq. formaldehyde (40.3 ml, 538 mmol). The solution was heated to 80 °C during which time a precipitate appeared and gradually redissolved over 1 hour. The reaction mixture was stirred at 80 °C overnight then cooled to room temperature. The solvents were removed under reduced pressure and the residue was dissolved in ethyl acetate (200 ml), washed successively with 1 M hydrochloric acid and brine, dried over anhydrous magnesium sulfate and filtered. The filtrate was concentrated to give the title compound as a clear oil (13.37 g, 97%). ¹H-NMR; δ (CDCl₃), 6.29 (1H, s), 5.65 (1H, s), 2.34-2.28 (2H, m), 1.54-1.26 (4H, m), 0.94 (3H, t, J = 7.1 Hz).

Step B: 4S-Benzyl-3-(2-butyl-acryloyl)-5,5-dimethyl-oxazolidin-2-one

2-Butyl acrylic acid (21.5 g, 168 mmol) was dissolved in dry THF (500 ml) and cooled to -78 °C under a blanket of argon. Triethylamine (30 ml, 218 mmol) and pivaloyl chloride (21 ml, 168 mmol) were added at such a rate that the temperature remained

below -60 °C. The mixture was stirred at -78 °C for 30 minutes, warmed to room temperature for 2 hours and finally cooled back to -78 °C.

In a separate flask, 4S-benzyl-5,5-dimethyl-oxazolidin-2-one was dissolved in dry THF (500ml) and cooled to -78 °C under a blanket of argon. n-Butyllithium (2.4 M solution in hexanes, 83 ml, 200 mmol) was added slowly and the mixture was stirred for 30 minutes at room temperature. The resulting anion was transferred *via* a cannula into the original reaction vessel. The mixture was allowed to warm to room temperature and was stirred overnight at room temperature. The reaction was quenched with 1 M potassium hydrogen carbonate (200 ml) and the solvents were removed under reduced pressure. The residue was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over anhydrous magnesium sulphate, filtered and concentrated under reduced pressure to give an orange oil. TLC analysis revealed the presence of unreacted chiral auxiliary in addition to the required product.

A portion of the material (30 g) was dissolved in dichloromethane and flushed through a silica pad to give pure title compound as a yellow oil (25.3 g). ¹H-NMR; δ (CDCl₃), 7.31-7.19 (5H, m), 5.41 (2H, s), 4.51 (1H, dd, J = 9.7 & 4.2 Hz), 3.32 (1H, dd, J = 14.2 & 4.2 Hz), 2.82 (1H, dd, J = 14.2 & 9.7 Hz), 2.40-2.34 (2H, m), 1.48-1.32 (4H, m), 1.43 (3H, s), 1.27 (3H, s), 0.91 (3H, t, J = 7.1 Hz). Some chiral auxiliary was recovered by flushing the silica pad with methanol.

Step C: 4S-Benzyl-3-[2-(benzyloxyamino-methyl)-hexanoyl]-5,5-dimethyl-oxazolidin-2-one (p-toluenesulfonic acid salt)

4S-Benzyl-3-(2-butyl-acryloyl)-5,5-dimethyl-oxazolidin-2-one (19.8 g, 62.8 mmol) was mixed with O-benzylhydroxylamine (15.4 g, 126 mmol) and stirred overnight at room temperature. The mixture was dissolved in ethyl acetate and the solution was washed with 1 M hydrochloric acid, 1 M sodium carbonate and brine, dried over anhydrous magnesium sulfate and filtered. The filtrate was concentrated under reduced pressure to afford a pale yellow oil (25.3 g) which was shown by NMR and HPLC analysis to contain 4S-benzyl-3-[2-(benzyloxyamino-methyl)-hexanoyl]-5,5-dimethyl-oxazolidin-2-

one (ca. 82% d.e.) along with a trace of starting material. The product was combined with another batch (26.9g, 76% d.e.) and dissolved in ethyl acetate (200 ml). p-Toluenesulfonic acid (22.7 g, 119 mmol) was added and the mixture was cooled to 0 °C. The title compound was obtained as a white crystalline solid by seeding and scratching. Yield: 25.2g, (34%, single diastereoisomer). A second crop (14.7 g, 20%, single diastereoisomer) was also obtained. ¹H-NMR;δ (CDCl₃), 7.89 (2H, d, J = 8.2 Hz), 7.37-7.12 (10H, m), 7.02 (2H, d, J = 6.9 Hz), 5.28-5.19 (2H, m), 4.55 (1H, m), 4.23 (1H, m), 3.93 (1H, m), 3.58 (1H, m), 2.58 (1H, m), 2.35 (3H, s), 1.67-1.51 (2H, m), 1.29-1.16 (4H, m), 1.25 (3H, s), 1.11 (3H, s), 0.80-0.75 (3H, m).

Step D: 2R-(Benzyloxyamino-methyl)-hexanoic acid

4S-Benzyl-3-[2R-(benzyloxyamino-methyl)-hexanoyl]-5,5-dimethyl-oxazolidin-2-one p-toluenesulfonic acid salt (25.2 g, 40.2 mmol) was partitioned between ethyl acetate and 1 M sodium carbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and evaporated under reduced pressure. The residual oil was dissolved in THF (150 ml) and water (50 ml), cooled to 0 °C and treated with lithium hydroxide (1.86 g, 44.2 mmol). The solution was stirred for 30 minutes at 0 °C, then overnight at room temperature. The reaction was acidified to pH4 with 1 M citric acid and the solvents were removed. The residue was partitioned between dichloromethane and 1 M sodium carbonate. The basic aqueous layer was acidified to pH4 with 1M citric acid and extracted three times with ethyl acetate. The combined organic layers were dried over anhydrous magnesium sulfate, filtered and concentrated to provide the title compound as a colourless oil (7.4 g, 73%). ¹H-NMR;δ (CDCl₃), 8.42 (2H, br s), 7.34-7.25 (5H, m), 4.76-4.66 (2H, m), 3.20-3.01 (2H, m), 2.73 (1H, m), 1.70-1.44 (2H, m), 1.34-1.22 (4H, m) and 0.92-0.86 (3H, m).

WO 01/10835

PCT/GB99/02629

24

Step E: 2R-[(Benzyloxy-formylamino)-methyl]-hexanoic acid

To a solution of 2R-(Benzyloxyamino-methyl)-hexanoic acid (30.6 g, 0.12 mol) in dry THF (300 ml) was added formic acetic anhydride (26.8 ml, 0.31 mol) at 0 °C. Triethylamine (18.5 ml, 0.13 mol) was added and the reaction was stirred for 1 h at 0 °C and 60 h at room temperature. The solvent was removed *in vacuo* to yield the title compound as a yellow oil (33.6 g, 99%) which was used in Step F without further purification. ¹H-NMR; (CDCl₃, rotamers), 8.20-8.08 (0.7H, br s), 8.07-7.92 (0.3H, br s), 7.50-7.25 (5H, br m), 5.07-4.70 (2H, br m), 3.95-3.52 (2H, br m), 2.90-2.66 (1H, br s), 1.72-1.20 (6H, br m), 1.00-0.78 (3H, br s). LRMS: +ve ion 280 [M+1].

Step F: 2R-[(Benzyloxy-formyl-amino)-methyl]-hexanoic acid pentafluorophenyl ester

To a solution of 2R-[(Benzyloxy-formylamino)-methyl]-hexanoic acid (7.8 g, 19.9 mmol) in dry THF (500 ml) was added pentafluorophenol (44.3 g, 0.24 mol), EDC (27.7 g, 0.14 mol) and HOBt (16.2 g, 0.12 mol). The reaction was stirred overnight at room temperature. The solvent was removed *in vacuo* and the residue was dissolved in ethyl acetate, washed successively with 1 M sodium carbonate (3 x 500 ml) and water (1 x 500 ml), dried over anhydrous magnesium sulfate and filtered. The solvent was removed *in vacuo* to yield a yellow oil (60 g) that was purified by flash chromatography (5:1, hexane:ethyl acetate → 1:2 hexane:ethyl acetate) to yield a clear oil (42.0 g, 79%). ¹H-NMR; δ(CDCl₃, rotamers), 8.20-8.09 (0.7H, br s), 8.09-7.92 (0.3H, br s), 7.60-7.21 (5H, br m), 5.00-4.70 (2H, br m), 4.04-3.72 (2H, br m), 3.18-3.00 (1H, br s), 1.85-1.57 (2H, br m), 1.50-1.26 (4H, br m), 1.00-0.82 (3H, br m); LRMS: 466 [M+H].

Step G: Generic experimental procedure for the synthesis of an array of amides








The coupling of amines to 2R-[(Benzyloxy-formyl-amino)-methyl]-hexanoic acid pentafluorophenyl ester was carried out on a Zymate XPii laboratory robot. To solutions of the pentafluorophenol ester (55.8 mg, 0.12 mmol) in dichloromethane (2





ml) were added individual amines (0.25 mmol) and the reaction mixtures were stirred at RT for 60 h. Purification was effected by removing excess amine and pentafluorophenol using scavenger resins. The pentafluorophenol was removed using a three fold excess (0.36 mmol) of A-26 carbonate resin (3.5 mmol loading). The resin was added to the reaction mixtures and agitated for 24 h, after which time it was filtered off. The excess amines were removed using a three-fold excess (0.36 mmol) of methylisocyanate polystyrene resin (1.2 mmol loading). The resin was added to the reaction mixtures and agitated for 4 h, after which time it was filtered off. The solvent was removed *in vacuo* using a Savant Speed Vac Plus to yield the coupled products. Yields were not calculated and the purity and integrity of each compound was verified using HPLC and LRMS.

Step H: Generic transfer hydrogenation procedure

Products from Step G were individually taken up in an ethanol (2.7ml) and cyclohexene (0.3 ml), 20% palladium on charcoal was added and the reactions stirred at 80 °C for 24 h. The Pd/C was filtered off and the solvent was removed *in vacuo* using a Savant Speed Vac Plus to yield the title compounds (examples **2-12**, Table 1). Yields were not calculated and the purity and integrity of each compound were verified using HPLC and LRMS


Table 1

Example	Structure	Mass Spectral Data	HPLC	Purification
2		347 (M+1, 100)	RT 18.5 min 100%	Resins
3		271 (M+1, 100), 293 (M+Na, 50)	RT 19.4 min 100%	Resins
4		257 (M+1, 50)	RT 24.4 min 100%	Resins
5		258 (M+1, 100)	RT 3.1 min and 3.5 min	Resins, Prep HPLC
6		258 (M+1, 100)	RT 4.0 min	Resins, Prep HPLC
7		300 (M+1, 100)	RT 4.2 min and 4.7 min (TFA salt)	Resins, Prep HPLC
8		271 (M+1, 100)	RT 18.5 min	Resins

9		287 (M+1, 100)	RT 3.0 min and 3.4 min	Resins, Prep HPLC
10		295 (M+1, 100)	Only prep RT	Prep HPLC
11		351 (M+Na, 100)	RT 7.6 min (grad 220nm)	Ion exchange Prep HPLC
12		330 (M+1, 100), 351 (M+Na, 50)	RT 16.8 min 100%	Resins


Example 13

**2R,4-{2-[(Formyl-hydroxy-amino)-methyl]-hexanoyl}-1,1-dimethyl-
piperazinium iodide**



The title compound was prepared using the same procedure as for Examples 2 to 12, except for the final methylation (see Scheme 4)

Scheme 4



Reagents and conditions: G. *N*-methylpiperazine; H. H₂, Pd/C, EtOH; I. MeI, dry THF.


Step I: 2R,4-{2-[(Formyl-hydroxy-amino)-methyl]-hexanoyl}-1,1-dimethyl-piperazinium iodide

To a solution of *N*-hydroxy-*N*-[2R-(4-methyl-piperazine-1-carbonyl)-hexyl]-formamide (46 mg, 0.17 mmol) in anhydrous THF (5 ml) was added methyl iodide (22 μ l, 0.34 mmol) dropwise at 0 °C. The reaction mixture was stirred at 0 °C for 1 h and at room temperature for 18 h. The solvent was removed *in vacuo* to yield the title compound as a white hygroscopic solid (68 mg, 97%). ¹H-NMR; CD₃OD, rotamers), 8.31 (0.7H, s), 7.88 (0.3H, s), 4.44-3.20 (17H, m), 1.75-1.20 (6H, m), 1.00-0.87 (3H, t, J = 6.6 Hz). LRMS: +ve ion 286 [M].

The compounds of Examples 14-17 were prepared from 2R-[(benzyloxy-formyl-amino)-methyl]-hexyl pentafluorophenyl ester (Example 2) and the appropriate L-tert-leucine derivatives by analogy with the method described in Example 2.

Example 14


2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid [2,2-dimethyl-1S-(piperidine-1-carbonyl)-propyl]-amide



White foam. LRMS: +ve ion 392 [M+Na], -ve ion 368 [M-H]. HPLC: RT = 20.7min.
(Purity 88%)

Example 15


**2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid
[1S-(3,4-dihydro-1H-isoquinoline-2-carbonyl)-2,2-dimethyl-propyl]-amide**



White foam. LRMS: +ve ion 440 [M+Na], -ve ion 416 [M-H]. HPLC: RT = 20.7min.
(Purity 91%)

Example 16


2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-4-hydroxy-piperidine-1-carbonyl)-2,2-dimethyl-propyl]-amide



White foam. LRMS: +ve ion 498 [M+Na], -ve ion 474 [M-H]. HPLC: RT = 21.0 min.
(Purity 96%).


Example 17

2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-piperazine-1-carbonyl)-2,2-dimethyl-propyl]-amide



White foam. LRMS: +ve ion 461 [M+H]. HPLC: RT = 16.6min. (Purity 86%).


32



White solid. ¹H-NMR; δ (CDCl₃, rotamers), 8.36 (0.3H, s), 7.79 (0.7H, s), 7.23 (6H, m), 5.30 (1H, m), 5.09 (2H, m), 3.96 (0.3H, dd, J = 14.2 & 8.6 Hz), 3.71 (0.7H, dd, J = 13.9 & 10.1 Hz), 3.47 (1H, m), 3.09 (1H, s), 3.06 (2H, s), 2.92 (1H, s), 2.91 (2H, s), 2.82 (3H, m), 1.68 (1H, m), 1.33 (5H, m) and 0.86 (3H, m). ¹³C-NMR; δ (CDCl₃, rotamers), 175.0, 173.1, 171.0, 170.7, 135.9, 129.0, 128.9, 128.8, 67.6, 67.3, 52.5, 49.2, 46.7, 46.4, 46.1, 45.9, 45.1, 37.7, 37.5, 37.4, 36.5, 36.4, 30.0, 29.8, 22.9 and 14.3. LRMS: +ve ion 444 [M+Na], 422 [M+H].

Example 20


4S-Dimethylcarbamoyl-4-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoylamino}-butyric acid benzyl ester



Pale yellow oil. LRMS: +ve ion 458 [M+Na], -ve ion 434 [M-H].

Example 21


(5S-Dimethylcarbamoyl-5-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoylamino}-pentyl)-dimethyl-ammonium chloride



Yellow oil. $^1\text{H-NMR}$; δ (CDCl_3), 7.77 (1H, s), 7.45 (1H, d, $J = 8.9$ Hz), 4.99 (1H, m), 3.81 (1H, m), 3.46 (1H, m), 3.09 (6H, s), 2.98 (3H, m), 2.97 (3H, s), 2.95 (3H, s), 1.51 (12H, m) and 0.88 (3H, m). $^{13}\text{C-NMR}$; δ (CDCl_3), 173.6, 171.5, 158.8, 58.2, 53.6, 48.6, 45.4, 37.6, 36.2, 31.4, 30.2, 29.7, 24.7, 23.0, 22.5 and 14.3. LRMS: +ve ion 373 $[\text{M}+\text{H}]$.


Example 22

2R-[(Formyl-hydroxy-amino)-methyl]-butyric acid (1S-carbamoyl-2,2-dimethyl-propyl) amide.



¹³C-NMR; δ (CDCl₃), 173.1, 55.5, 54.9, 51.7, 48.4, 48.0, 46.6, 38.9, 38.8, 36.3, 36.1, 31.3, 27.0, 26.9, 23.9, 23.8 and 12.1. LRMS: +ve ion 324 [M+Na] 300 [M-H].


2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid (1S-carbamoyl-2,2-dimethyl-propyl) amide.



SUBSTITUTE SHEET (RULE 26)

Example 24


2R-[Formyl-hydroxy-amino)-methyl]-hexanoic acid (1S-dimethyl-carbamoyl-4-guanidinobutyl)-amide



White powder. $^1\text{H-NMR}$; $\delta(\text{CD}_3\text{OD}$, rotamers), 8.12 (0.1H, s), 7.60 (0.9H, s), 4.90 (1H, m), 3.67 (1H, dd, $J = 12.2$, 12.2 Hz), 3.38 (1H, m), 3.22-3.09 (2H, m), 3.11 (3H, s), 3.02 (1H, m), 2.94 (3H, s), 1.74-1.47 (5H, m), 1.47-1.20 (5H, m) and 0.90 (3H, t, $J = 6.6$ Hz). $^{13}\text{C-NMR}$; $\delta(\text{CD}_3\text{OD}$, rotamers), 174.4, 172.0, 157.9, 55.9, 49.0, 45.0, 41.4, 37.7, 36.2, 30.6, 29.8, 29.7, 25.1, 23.2 and 14.4.
LRMS: +ve ion 373 [M+H].

Example 25

[2R-(4-chlorophenyl)-3-(formyl-hydroxy-amino)-propionylamino]-2S-3,3,N,N-tetramethyl-butamide



WO 01/10835

PCT/GB99/02629

37

White solid. ¹H-NMR: δ(CD₃OD, rotamers), 8.24 (0.3H, s), 8.10 (1H, br m), 7.84 (0.7H, s), 7.03 (2H, d, J = 8 Hz), 6.70 (2H, d, J = 7 Hz), 3.68 (1H, m), 3.35 (3H, m), 2.70 (3H, m), 1.65-1.10 (6H, m) and 0.90 (3H, t, J = 7.0 Hz); ¹³C-NMR: δ(CD₃OD, rotamers), 176.5, 176.1, 157.3, 131.6, 131.2, 116.7, 53.9, 46.1, 45.1, 42.9, 36.1, 31.7, 31.2, 24.1 and 14.7.; LRMS: +ve ion 309 [M+H], 331 [M+Na]; -ve ion 307 [M-H].

Biological Example A

Demonstration of antibacterial effect

a).

Minimal inhibitory concentrations (MIC) of inhibitors against *E. coli* strain DH5α (Genotype; F-φ80d*lacZ*ΔM15Δ(*lacZYA-argF*)U169 *deoR recA1 endA1 hsdR17*(r_k⁻, m_k⁺)*phoA supE44*λ⁻ *thi-1 gyrA96 relA1*) obtained from GibcoBRL Life Technologies, *Enterobacter cloacae* (American Type Culture Collection number 13047), *Klebsiella pneumoniae* (American Type Culture Collection number 13883) or *Staphylococcus capitis* (American Type Culture Collection number 35661) were determined as follows. Stock solutions of test compound (Compounds 1 and 2 from Examples 1 and 2 respectively (both isomer A)) and three standard laboratory antibiotics, carbenicillin (Sigma, catalogue No. C3416), kanamycin (Sigma, catalogue No. K4000) and chloramphenicol (Sigma, catalogue No. C1919), were prepared by dissolution of each compound in dimethylsulfoxide at 10mM. For the determination of the minimal inhibitory concentration, two fold serial dilutions were prepared in 2xYT broth (typtone 16g/1, yeast extract 10g/1, sodium chloride 5g/1 obtained from BIO 101 Inc, 1070 Joshua Way, Vista, CA92083, USA) to yield 0.05 ml compound-containing medium per well. Inocula were prepared from cultures grown overnight in 2xYT broth at 37°C. Cell densities were adjusted to absorbance at 660nm (A₆₆₀) = 0.1; the optical density-standardised preparations were diluted 1:1000 in 2xYT broth; and each well inoculated with 0.05ml of the diluted bacteria. Microtitre plates were incubated at 37°C for 18 hours in a humidified incubator. The MIC (μM) was recorded as the lowest drug concentration that inhibited visible growth. The compounds of the Examples inhibited bacterial growth. For example,

the compound of Example 7 had an MIC against *E. coli* of 12.5 μ M.

Biological Example B

i) *Cloning of the Escherichia coli PDF gene.*

The *E. coli* PDF gene was cloned in pET24a(+) (designated pET24-PDF) and was used to transform BL21 DE3 cells from Novagen Inc, (Madison, Wisconsin).

Clones were selected at 37°C on YT agar plates (8g/l typtone, 5g/yeast extract, NaCl 5g/l, agar 15g/l) supplemented with 30 μ g/ml kanamycin.

ii) *Expression of PDF*

A 20ml overnight culture of BL21 DE3 cells harbouring pET24-PDF was used to infect 500ml 2xYT broth (16g/l typtone, 10g/l yeast extract, NaCl 5g/l) containing 30ug/ml kanamycin in a 2 litre baffled flask and grown at 37°C with shaking to an OD₆₀₀ 0.6. The culture was then induced by adjusting the medium to 1.0mM isopropyl β -D thiogalactopyranoside (IPTG). The induction was allowed to proceed for a further 3 hours at 37°C, the cells were harvested by centrifugation and the cell pellet washed with 250ml phosphate buffered saline (PBS) and the pellet stored at -70°C.

iii) *Preparation of soluble protein fraction.*

The cells from a 1 litre expression were resuspended in 2x 25ml of ice cold phosphate buffered saline. The cell suspension was sonicated on ice using an MSE Soniprep 150 fitted with a medium probe and at an amplitude of 20-25 microns in 6x20 second pluses. The resulting suspension was then cleared by centrifugation at 20,000 xg for 15 minutes. The supernatant was then used for further purification of the enzyme.

iv) *PDF Purification*

E. coli lysate from a 1l culture in phosphate buffered saline (PBS) were adjusted to 2M ammonium sulphate. A 15ml phenyl sepharose column was equilibrated with PBS/2M ammonium sulphate at 4°C. The lysate was loaded on the column and washed with equilibration buffer. The column was eluted by reducing the ammonium sulphate concentration from 2M to 0M over 10 column volumes. 5ml fractions were collected and analysed by SDS-PAGE. The fractions containing the majority of the 20kDa PDF were pooled. The pooled fractions were concentrated using a 3kDa cutoff membrane to a volume of 5ml. The fraction was then loaded onto a Superdex 75 (size exclusion chromatography) column equilibrated in PBS. The concentrated PDF pool eluted at one ml/min at 4°C and 5ml fractions collected and analysed by SDS-PAGE. The purest fractions were pooled and stored at -70°C.

(v) *PDF in vitro assay*

The assay was performed in a single 96 well plate in a final volume of 100µl containing:

- 20µl PDF (4µg/ml)
- 20µl 100mM Hepes pH 7.0 + 1M KCl + 0.05% Brij
- 10µl serial dilution of test compound in 20% DMSO
- 50µl formyl-Met-Ala-Ser (8mM)

The assay was incubated at 37°C for 30 minutes. The free amino group of the deformylated (Met-Ala-Ser) product was detected using fluorescamine, by the following additions:


- 50µl 0.2M borate pH 9.5
- 50µl fluorescamine (150µg/ml in dry dioxane)

Fluorescence was quantified on SLT Fluostar plate reader using an excitation wavelength of 390nm and an emission wavelength of 485nm. Standard control reactions are a no inhibitor reaction which provides the zero inhibition figure and a no enzyme and no inhibitor reaction which provides the 100% inhibition figure. The data was analysed by conversion of the fluorescence units to % inhibition and the inhibitor concentration plotted against % inhibition. The data was fitted to a sigmoidal function : $y = A + ((B - A) / (1 + ((C/x)^D)))$, wherein A represents zero inhibition, B represents 100% inhibition and C represents the IC_{50} , D represents the slope. The IC_{50} represents the concentration of inhibitor (nM) required to decrease enzyme activity by 50%.



The compounds of the invention were found to inhibit bacterial PDF *in vitro*.

Claims :

1. The use of a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt hydrate or solvate thereof in the preparation of an antibacterial composition:



wherein R_2 represents a substituted or unsubstituted $\text{C}_1\text{-C}_6$ alkyl, cycloalkyl($\text{C}_1\text{-C}_6$ alkyl)-, or aryl($\text{C}_1\text{-C}_6$ alkyl)- group, and A represents a group of formula (IA), or (IB):



wherein R_4 represents the side chain of a natural or non-natural alpha amino acid, and R_5 and R_6 are each independently hydrogen or $\text{C}_1\text{-C}_6$ alkyl, heterocyclic or aryl($\text{C}_1\text{-C}_6$ alkyl)-, or R_5 and R_6 when taken together with the nitrogen atom to which they are attached form an optionally substituted saturated heterocyclic ring of 3 to 8 atoms which ring is optionally fused to a carbocyclic or second heterocyclic ring,

characterised in that the said compound is selected from the group consisting of

N-[3S-(4-benzylpiperidine-1-carbonyl)-2,2-dimethyl-propyl]-3-cyclopentyl-2R-[(formyl-hydroxy-amino)-methyl]-propionamide,

N-[2R-(4-benzyl-piperidine-1-carbonyl)-hexyl]-N-hydroxy-formamide,

N-hydroxy-N-[2R-(2-methyl-piperidine-1-carbonyl)-hexyl]-formamide,

N-hydroxy-N-[2R-(piperidine-1-carbonyl)-hexyl]-formamide,

N-hydroxy-N-[2R-(piperazine-1-carbonyl)-hexyl]-formamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid pyrrolidin-1-ylamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid methyl-(1-methyl-piperidin-4-yl)-amide,

N-[2R-(azepane-1-carbonyl)-hexyl]-N-hydroxy-formamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (4-methyl-piperazin-1-yl)-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid diisopropylamide,

1-[2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl]-piperidine-3-carboxylic acid ethyl ester,

4-[2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl]-piperazine-1-carboxylic acid ethyl ester,

4-[2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl]-1,1-dimethyl-piperazinium iodide,

2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid [2,2-dimethyl-1S-(piperidine-1-carbonyl)-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(3,4-dihydro-1*H*-isoquinoline-2-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-4-hydroxy- piperidine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-piperazine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (3-benzylsulfanyl-1S- dimethylcarbamoyl-propyl)-amide,

3S-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoylamino}-*N,N*-dimethyl-succinamic acid benzyl ester,

4S-dimethylcarbamoyl-4-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl-amino}-butyric acid benzyl ester,

(5S-dimethylcarbamoyl-5-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl-amino}-pentyl)-dimethyl-ammonium chloride,

2R-[(formyl-hydroxy-amino)-methyl]-butyric acid (1-carbamoyl-2,2-dimethyl-propyl) amide,

2-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (1-carbamoyl-2,2-dimethyl-propyl) amide,

2R-[formyl-hydroxy-amino)-methyl]-hexanoic acid (1-dimethyl-carbamoyl-4-guanidinobutyl)-amide,

2R-[2-(4-chlorophenyl)-3-(formyl-hydroxy-amino)-propionylamino]-2S-3,3,*N,N*-tetramethyl-butylamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [2(3,4-dihydroxy-phenyl)-ethyl]-amide,


2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [2(4-hydroxyphenyl)-ethyl]-amide,

and pharmaceutically and veterinarily acceptable salts, hydrates and solvates thereof.


2. A method for the treatment of bacterial infections in humans and non-human mammals, which comprises administering to a subject suffering such infection an antibacterially effective dose of a compound as specified in claim 1.

3. A method for the treatment of bacterial contamination by applying an antibacterially effective amount of a compound as specified in claim 1 to the site of contamination;

4. A compound of formula (I) or a pharmaceutically or veterinarily acceptable salt hydrate or solvate thereof



wherein R_2 represents a substituted or unsubstituted $\text{C}_1\text{-C}_6$ alkyl, cycloalkyl($\text{C}_1\text{-C}_6$ alkyl)-, or aryl($\text{C}_1\text{-C}_6$ alkyl)- group, and A represents a group of formula (IA), or (IB):



characterised in that the said compound is selected from the group consisting of

N-[2R-(4-benzyl-piperidine-1-carbonyl)-hexyl]-N-hydroxy-formamide,

N-hydroxy-N-[2R-(2-methyl-piperidine-1-carbonyl)-hexyl]-formamide.

N-hydroxy-N-[2R-(piperidine-1-carbonyl)-hexyl]-formamide,

N-hydroxy-N-[2R-(piperazine-1-carbonyl)-hexyl]-formamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid pyrrolidin-1-ylamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid methyl-
(1-methyl- piperidin-4-yl)-amide,

N-[2R-(azepane-1-carbonyl)-hexyl]-N-hydroxy-formamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (4-methyl-
piperazin-1-yl)- amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid diisopropylamide,

1-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl}-piperidine-3-carboxylic acid ethyl ester,

4-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl}-piperazine-1-carboxylic acid ethyl ester,

4-{2R-[(formyl-hydroxy-amino)-methyl]-hexanoyl}-1,1-dimethyl-piperazinium iodide,

2R-[(Formyl-hydroxy-amino)-methyl]-hexanoic acid [2,2-dimethyl-1S-(piperidine-1-carbonyl)-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(3,4-dihydro-1*H*-isoquinoline-2-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-4-hydroxy-piperidine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [1S-(4-benzyl-piperazine-1-carbonyl)-2,2-dimethyl-propyl]-amide,

2-[(formyl-hydroxy-amino)-methyl]-hexanoic acid (1-carbamoyl-2,2-dimethyl-propyl) amide,

2R-[2-(4-chlorophenyl)-3-(formyl-hydroxy-amino)-propionylamino]-2S-3,3,*N,N*-tetramethyl-butylamide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [2(3,4-dihydroxy-phenyl)-ethyl]-amide,

2R-[(formyl-hydroxy-amino)-methyl]-hexanoic acid [2(4-hydroxyphenyl)-ethyl]-amide,

and pharmaceutically and veterinarily acceptable salts, hydrates and solvates thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 February 2001 (15.02.2001)

PCT

(10) International Publication Number
WO 01/10835 A1

(51) International Patent Classification⁷: **C07D 211/16**,
295/18, A61K 31/16, C07D 295/22, C07C 259/06, C07D
211/60, 295/20, 217/06, 211/48, C07C 321/16, 279/14,
A61K 31/445, 31/495, 31/47, A61P 31/04, C07D 211/58

(21) International Application Number: PCT/GB99/02629

(22) International Filing Date: 10 August 1999 (10.08.1999)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US):
**BRITISH BIOTECH PHARMACEUTICALS LIM-
ITED** [GB/GB]; Watlington Road, Cowley, Oxford OX4
5LY (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HUNTER, Michael,
George** [GB/GB]; British Biotech Pharmaceuticals Lim-
ited, Watlington Road, Cowley, Oxford OX4 5LY (GB).
BECKETT, Raymond, Paul [GB/GB]; British Biotech
Pharmaceuticals Limited, Watlington Road, Cowley,
Oxford OX4 5LY (GB). **CLEMENTS, John, Martin**
[GB/GB]; British Biotech Pharmaceuticals Limited,
Watlington Road, Cowley, Oxford OX4 5LY (GB).
WHITTAKER, Mark [GB/GB]; British Biotech Phar-
maceuticals Limited, Watlington Road, Cowley, Oxford

OX4 5LY (GB). **DAVIES, Stephen, John** [GB/GB];
British Biotech Pharmaceuticals Limited, Watlington
Road, Cowley, Oxford OX4 5LY (GB). **PRATT, Lisa,
Marie** [GB/GB]; British Biotech Pharmaceuticals Lim-
ited, Watlington Road, Cowley, Oxford OX4 5LY (GB).
SPAVOLD, Zoe, Marie [GB/GB]; British Biotech Phar-
maceuticals Limited, Watlington Road, Cowley, Oxford
OX4 5LY (GB). **LAUNCHBURY, Steven** [GB/GB];
British Biotech Pharmaceuticals Limited, Watlington
Road, Cowley, Oxford OX4 5LY (GB).

(74) Agent: **WALLS, Alan, J.**; British Biotech Pharmaceu-
ticals Limited, Watlington Road, Cowley, Oxford OX4 5LY
(GB).

(81) Designated States (national): AU, BR, CA, CN, CZ, GB,
HU, IL, JP, KR, MX, NO, NZ, PL, RU, SG, SK, TR, UA,
US, ZA.


(84) Designated States (regional): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

Published:


— With international search report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.


(54) Title: ANTIBACTERIAL AGENTS



(I)



(IA)



(IB)

(57) Abstract: Selected compounds of formula (I) are antibacterial agents: formula (I) wherein R₂ represents a substituted or un-substituted C₁-C₆ alkyl, cycloalkyl (C₁-C₆ alkyl)- or aryl (C₁-C₆ alkyl)- group, and A represents a group of formula (IA), or (IB) wherein R₄ represents the side chain of a natural or non-natural alpha amino acid, and R₅ and R₆ are each independently hydrogen or C₁-C₆ alkyl, heterocyclic or aryl (C₁-C₆ alkyl)-, R₅ and R₆ when taken together with the nitrogen atom to which they are attached from an optionally substituted saturated heterocyclic ring of 3 to 8 atoms which ring is optionally fused to a carbocyclic or second heterocyclic ring.

WO 01/10835 A1

Banner & Witcoff Ref. No. 010180.00012
Client Ref. No. 205/AJW

JOINT DECLARATION FOR PATENT APPLICATION

As the below named inventors, we hereby declare that:

Our residence, post office address and citizenship are as stated below next to our names;

We believe we are the original, first and joint inventors of the subject matter which is claimed and for which a patent is sought on the invention entitled ANTIBACTERIAL AGENTS, the specification of which

- ☐ is attached hereto.
☒ was filed on February 11, 2002 as Application Serial Number 10/049,274 and was amended on _____ (if applicable).
☒ was filed under the Patent Cooperation Treaty (PCT) and accorded International Application No. PCT/GB99/02629, filed August 10, 1999, and amended on _____ (if any).

We hereby state that we have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We hereby acknowledge the duty to disclose information which is material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56(a).

Prior Foreign Application(s)

We hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application(s) for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Country	Application No.	Date of Filing (day month year)	Date of Issue (day month year)	Priority Claimed Under 35 U.S.C. §119

Prior United States Provisional Application(s)

We hereby claim priority benefits under Title 35, United States Code, §119(e)(1) of any U.S. provisional application listed below:

U.S. Provisional Application No.	Date of Filing (day month year)	Priority Claimed Under 35 U.S.C. §119(e)(1)

Prior United States Application(s)

We hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Date of Filing (Day, Month, Year)	Status — Patented, Pending, Abandoned

Banner & Witcoff Ref. No. 010180.00012
Client Ref. No. 205/AJW

Power of Attorney

And we hereby appoint, both jointly and severally, as our attorneys with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith the practitioners at:

Customer Number: 22907 (WDC)

Please address all correspondence and telephone communications to the address and telephone number for this Customer Number.

We hereby declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1-00
Signature M.G. Hunter Date 16th July 2002
Full Name of First Inventor Hunter Michael George
Family Name First Given Name Second Given Name
Residence United Kingdom CBX Citizenship British
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB

2-00
Signature Paul Beckett Date 22nd July 2002
Full Name of Second Inventor Beckett Raymond Paul
Family Name First Given Name Second Given Name
Residence United Kingdom CBX Citizenship British
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB

3-00
Signature John Clements Date 22nd August 2002
Full Name of Third Inventor Clements John Martin
Family Name First Given Name Second Given Name
Residence United Kingdom CBX Citizenship British
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB

4-00
Signature M. Whittaker Date 30 July 2002
Full Name of Fourth Inventor WHITTAKER Mark
Family Name First Given Name Second Given Name
Residence United Kingdom CBX Citizenship British
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB

Signature Stephen Davies Date 29 July 2002
Full Name of Fifth Inventor Davies Stephen John
Family Name First Given Name Second Given Name
Residence United Kingdom GB Citizenship British
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB

Signature *Lisa Pratt* Date 13 August 2002
Full Name of Sixth Inventor Pratt Lisa Marie
Residence United Kingdom GBX Family Name First Given Name Second Given Name
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB
Citizenship British

Signature Zoe Spavold Date 17 August 2002
Full Name of Seventh Inventor Spavold Zoe Marie
Family Name First Given Name Second Given Name
Residence United Kingdom GB GB GB
Citizenship British British British
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB

Signature Steven Launchbury Date 2nd SEPT 2002
Full Name of Eighth Inventor Launchbury Steven
Family Name First Given Name Second Given Name
Residence United Kingdom GB Citizenship British
Post Office Address British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 6LY, GB